Paediatric Hypoglycaemia How to order on TRAKCARE

'PAEDS HYPOGLYCAEMIA - TIME CRITICAL SAMPLES'

(Samples required PRIOR to administration of glucose)

TO ORDER THE 'TIME CRITICAL SAMPLES' THEN FOLLOW THESE STEPS:

- 1. Ensure patient highlighted on ED screen then click on "New Request" tab.
- 2. CLICK ON "LABS-CHILD"
- 3. On the right side of screen type the requested test in section titled "item" then hit 'F6'key.
- 4. SELECT APPROPRIATE TEST IN POP-UP SCREEN

ALWAYS SELECT THE '- CHILD' OPTION FOR THE TEST REQUESTED (WHEN AVAILABLE)

HINTS:

- FOR FFAS TYPE 'FREE' THEN HIT F6 KEY
- FOR C PEPTIDE ENSURE THERE IS A SPACE EITHER SIDE OF THE '-' BEFORE HITTING F6
- FOR BETA OH BUTYRATE TYPE 'BETA' THEN HIT F6 KEY
- REMEMBER LACTATE IS TAKEN ON A CAPILLARY GAS TUBE AND PROCESSED IN THE ED.

'PAEDS HYPOGLYCAEMIA - ADDITIONAL ESSENTIAL'

(Sample can be taken **AFTER** administration of glucose)

TO ORDER THE 'ADDITIONAL ESSENTIAL' SAMPLES THEN FOLLOW THE SAME PROCESS AS ABOVE.

ALWAYS SELECT THE '- CHILD' OPTION FOR THE TEST REQUESTED (WHEN AVAILABLE)

HINTS:

- FOR ACYLCARNITINE ENSURE YOU SELECT THE OPTION WITH 'BS' AT THE END.
- FOR URINARY ORGANIC ACIDS TYPE 'ORGANIC' THEN HIT F6 KEY

ONLY STICK **ONE** 'TRAKCARE' LABEL TO THE CORRESPONDING TUBE.
THE REMAINING LABELS SHOULD REMAIN ON THE REQUEST FORM AND ACCOMPANY THE SAMPLES TO THE LAB